

Determination of a Wave Function Functional

Xiao-Yin Pan, Viraht Sahni, and Lou Massa

The Graduate School of the City University of New York, New York, New York 10016.

We propose expanding the space of variations in traditional variational calculations for the energy by considering the wave function ψ to be a functional of a set of functions $\chi : \psi = \psi[\chi]$, rather than a function. A constrained search in a subspace over all functions χ such that the functional $\psi[\chi]$ satisfies a sum rule or leads to a physical observable is then performed. An upper bound to the energy is subsequently obtained by variational minimization. The *rigorous* construction of such a constrained-search-variational wave function functional is demonstrated.

One of the mostly extensively employed and accurate approximation methods in quantum mechanics is the variational principle for the energy. Consider a quantum mechanical system with Hamiltonian operator $\hat{H}$. The ground state eigenenergies E and eigenfunctions Ψ for this system satisfy the Schrödinger equation $\hat{H}\Psi = E\Psi$. Next define the functional $I[\psi] = \int \psi^* \hat{H} \psi d\tau / \int \psi^* \psi d\tau$. Searching over all bounded, quadratically integrable functions ψ , one obtains $I[\Psi] = E$, with Ψ being the solution of the Schrödinger equation associated with energy E . Since $\delta I[\Psi] = 0$, the functional $I[\psi]$ is stationary for $\psi = \Psi$. In practice, an approximate function ψ of a particular analytical form is chosen to depend upon a number of variable parameters $c_i (i = 1, \dots, p)$. A least upper bound to the ground state energy E_0 is obtained by solving the p equations $\partial I[\psi] / \partial c_i = 0$, and employing the derived set of values of the parameters c_i to calculate $I[\psi]$. In application of the variational principle, however, the space of variations is limited by the choice of form of the function chosen for the approximate wave function. For example, if Gaussian or Slater-type orbitals are employed, then the variational space is limited to such functions. In this paper we propose the idea of overcoming this limitation by expanding the space over which the variations are performed. This then allows for a greater flexibility for the structure of the approximate wave function. We demonstrate the idea of expansion of the variational space by example.

We expand the space of variations by considering the approximate wave function to be a functional of the set of functions $\chi : \psi = \psi[\chi]$, rather than a function. The space of variations is expanded because the functional $\psi[\chi]$ can be adjusted through the function χ to reproduce any well behaved function. However, this space of variations is still too large for practical purposes, and so we consider a *subset* of this space. In addition to the function ψ being of a particular analytical form and dependent on the variational parameters c_i , the functions χ are chosen such that the functional $\psi[\chi]$ satisfies a constraint. Examples of such constraints on the wave function functional $\psi[\chi]$ are those of normalization or the satisfaction of the Fermi-Coulomb hole charge sum rule, or the requirement that it lead to

observables such as the electron density, the diamagnetic susceptibility, nuclear magnetic constant or any other physical property of interest. A constrained-search over all functions χ such that $\psi[\chi]$ satisfies a particular condition is then performed. With the functional $\psi[\chi]$ thus determined, the functional $I[\psi[\chi]]$ is then minimized with respect to the parameters c_i . In this manner both a particular system property of interest as well as the energy are obtained accurately, the latter being a consequence of the variational principle. We refer to this way of determining an approximate wave function as the *constrained-search-variational* method.

As an example of the method we consider its application to the ground state of the Helium atom. In atomic units $e = \hbar = m = 1$, the non-relativistic Hamiltonian is

$$\hat{H} = (-1/2)\nabla_1^2 - (1/2)\nabla_2^2 - Z/r_1 - Z/r_2 + 1/r_{12}, \quad (1)$$

where $\mathbf{r}_1, \mathbf{r}_2$ are the coordinates of the two electrons, r_{12} is the distance between them, $Z = 2$ is the atomic number. In terms of the Hylleraas coordinates[1] $s = r_1 + r_2$, $t = r_1 - r_2$, $u = r_{12}$, we choose the ground state wave function functional to be of the general form

$$\psi[\chi] = \Phi(\alpha, s)[1 - f(s, t, u)], \quad (2)$$

with $\Phi(\alpha, s) = (\alpha^3/\pi) \exp(-\alpha s)$, $f(s, t, u) = e^{-qu}(1 + qu)[1 - \chi(q, s, t, u)(1 + u/2)]$, where α and q are variational parameters. Any two-electron wave function may be expressed in this most general form.

In our example, we consider χ to be a function only of the variable s : $\psi = \psi[\chi(q, s)]$. Then, this wave function satisfies the electron-electron cusp condition[2], which in integral form is $\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)|_{\mathbf{r}_1 \rightarrow \mathbf{r}_2} = \Psi(\mathbf{r}_2, \mathbf{r}_2, \dots, \mathbf{r}_N)(1 + \frac{r_{12}}{2}) + \mathbf{r}_{12} \cdot \mathbf{C}(\mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N)$, where $\mathbf{r}_{12} = \mathbf{r}_1 - \mathbf{r}_2$ and $\mathbf{C}(\mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N)$ is an undetermined vector. This is readily seen by Taylor expanding $\psi[\chi(q, s)]$ about $u = 0$. In a similar manner, the wave function $\psi[\chi(q, s)]$ satisfies the electron-nucleus cusp coalescence condition[2], which is $\Psi(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_N)|_{\mathbf{r} \rightarrow 0} = \Psi(0, \mathbf{r}_2, \dots, \mathbf{r}_N)(1 - Zr_{12}) + \mathbf{r} \cdot \mathbf{A}(\mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N)$ with $\mathbf{A}(\mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N)$ being an undetermined vector, for $\alpha = 2$. For arbitrary value of the variational parameter α , the electron-nucleus coalescence condition

is not satisfied. (The differential form of these cusp conditions obtained from the integral versions is $d\Psi_{sp.av}/dr|_{r=0} = \zeta\Psi|_{r=0}$, where $\Psi_{sp.av}$ is the spherical average of the wave function about the singularity, $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, $\mathbf{r}_1$ and $\mathbf{r}_2$ the positions of the two particles, and with $\zeta = 1/2$ for electron-electron coalescence and $\zeta = -Z$ for electron-nucleus coalescence.)

The constraint to be applied to the wave function functional $\psi[\chi]$ is the normalization condition:

$$\int \psi^*[\chi]\psi[\chi]d\tau = 1. \quad (3)$$

A pictorial representation of the space of variation in the above example relative to the standard variational method is given in Fig.1.

The next step is the constrained search over functions $\chi(q, s)$ for which the condition of normalization is satisfied. Substitution of our $\psi[\chi]$ into Eq.(3) leads to

$$2\pi^2 \int_0^\infty ds |\Phi(\alpha, s)|^2 g(s) = 0, \quad (4)$$

where

$$g(s) = \int_0^s u du \int_0^u dt (s^2 - t^2) [f^2(s, u) - 2f(s, u)]. \quad (5)$$

Next consider Eq.(4). If the parameter α were fixed at say the value for which the energy is minimized by the Hydrogenic pre-factor $\Phi(\alpha, s)$, i.e. $\alpha = 27/16$, then there exist *many* functions $g(s)$ for which the condition of Eq.(4) is satisfied. These solutions correspond to the subspace C as shown in Fig.1.

On the other hand, if the parameter α is variable, then the only way in which the condition of Eq.(4) can be satisfied is if

$$g(s) = 0. \quad (6)$$

The requirement of Eq.(6) is equivalent to the constrained search over the entire subspace C. Substitution of $f(s, u)$ into Eq.(6) in turn is equivalent to a quadratic equation for the function $\chi(q, s)$:

$$a(q, s)\chi(q, s)^2 + 2b(q, s)\chi(q, s) + c(q, s) = 0, \quad (7)$$

where

$$a(q, s) = \int_0^s (s^2 u^2 - u^4/3)(1 + u/2)^2 (1 + qu)^2 e^{-2qu} du, \quad (8)$$

$$b(q, s) = - \int_0^s (s^2 u^2 - u^4/3)(1 + u/2)(1 + qu) [e^{-2qu}(1 + qu) - e^{-qu}] du, \quad (9)$$




FIG. 1: In the figure, the area of box A represents the space of all normalized square-integrable functions. Functions such as Gaussian or Slater orbitals are represented by lines. The space A is also representative of the functionals $\psi[\chi(s, t, u)]$. The subspace area B represents all normalized wave function functionals $\psi[\chi(s, t)]$, and its subspace C that of all normalized functionals $\psi[\chi(s)]$.

$$c(q, s) = \int_0^s (s^2 u^2 - u^4/3)(1 + qu)[e^{-2qu}(1 + qu) - 2e^{-qu}] du. \quad (10)$$

The integrals for the coefficients a , b , and c are solvable analytically. *Solution of the quadratic equation is then equivalent to searching over the entire subspace C. Subspace C is comprised of only two points.* These two points corresponding to the two solutions $\chi_1(q, s)$ and $\chi_2(q, s)$ are such that the two wave functions $\psi[\chi_1]$ and $\psi[\chi_2]$ are normalized. Rigorous upper bounds to the ground state energy are then obtained by variational minimization of the functional $I[\psi[\chi]]$ with respect to the parameters α and q . The details of the derivation and analytical expressions for the wave function functionals are to be given elsewhere.

The ground state energies obtained from the wave function functionals $\psi[\chi_1]$ and $\psi[\chi_2]$ are given in Table I together with those due to the energy-minimized Hydrogenic pre-factor $\Phi(s)$ with $\alpha = 27/16$, the Hartree-Fock(HF) value[3], and the 'exact' value of Pekeris[4]. The functions $\chi_1(q, s)$ and $\chi_2(q, s)$ at the respective energy minimum are plotted in Fig.2. We also quote energy minimized values determined from the wave function functionals $\psi[\chi_3]$ and $\psi[\chi_4]$ obtained by solution of the quadratic equation with α fixed at $\alpha = 27/16$, and q treated as the only variational parameter. (The functions (χ_3, χ_4) differ minimally from those of (χ_1, χ_2) .) The satisfaction of the virial theorem and the expectation value of the operator $W = r_1 + r_2$ are also quoted.

We note the following points of interest: (i) The functions $\chi_1(q, s)$, $\chi_2(q, s)$ are very different: $\chi_1(q, s)$ is a positive monotonically decreasing function whereas

$\chi_2(q, s)$ is a negative monotonically increasing function. They, however, are of about the same magnitude. Thus, the functionals $\psi[\chi_1]$ and $\psi[\chi_2]$ are different from each other. Similarly, the functionals $\psi[\chi_3]$ and $\psi[\chi_4]$ differ. (ii) In spite of the wave function functionals $\psi[\chi_1(q, s)]$ and $\psi[\chi_2(q, s)]$ being different, the corresponding energies are essentially equivalent. (iii) Note that $E[\chi_2] < E[\chi_1] < E[\chi_4] < E[\chi_3]$. (iv) The improvement in the energies noted in (iii) is also reflected in the corresponding satisfaction of the virial theorem. (v) As the constrained search is over the normalized functionals of subspace C, *both* solutions of the quadratic equation Eq.(7) lead to meaningful energies.

Our results demonstrate the advantage of the concept of a wave function functional. (The purpose of the paper is not to determine the most accurate wave function for the He atom ground state.) The energies obtained via the various wave function functionals are to our knowledge the most accurate one- and two-parameter results in the literature. A 1.93% error of the Hydrogenic pre-factor is reduced to errors of 0.45% – 0.43% for the four wave function functionals. The corresponding satisfaction of the virial theorem is 1.61% – 0.08%. As a further point of comparison, we note that our results are superior to those of HF, and the results of $\psi[\chi_1]$, $\psi[\chi_2]$ and $\psi[\chi_4]$ are also superior to a 3-parameter wave function calculation(CK) [5]. (This wave function is of the form of Eq.(2) with $f(s, u) = e^{-qu}(1 + qu)[1 - \beta e^{-2\mu s}(1 + u/2)]$ with $\alpha = 27/16$, $q = 0.562326$, $\mu = 0.099947$, $\beta = 0.88066$). The exact satisfaction of the virial condition by the pre-factor is a consequence of scaling, and that of HF due to the self-consistent nature of the solution. That of Pekeris is a consequence of the accuracy of the wave function. The expectation values of $W = r_1 + r_2$ are of course only expected to be as accurate as that of wave function, and not correct to second order as is the energy. The values obtained by the functionals $\psi[\chi_1]$ and $\psi[\chi_2]$ are an improvement over those of the pre-factor, reducing an error of 4.3% to 2.9%. They are also significantly superior to those obtained by the 3-parameter wave function which is in error by 3.9%. Similar improvements are observed for the expectations of other single particle operators. On the other hand, the high accuracy of the HF value is because in this theory single-particle expectation values are correct to second order [6, 9]. Our results also demonstrate that by expanding the space of variations, more accurate results for both the energy as well as other properties can be obtained with fewer variational parameters.

An improvement over the present results can be achieved as follows. (i) Expand the space of variations by considering χ to be a function of the variables s and t , or expand the space still further by considering χ to be

FIG. 2: The functions $\chi_1(q, s)$ and $\chi_2(q, s)$.

a function of the variables s, t , and u . Such an expansion of the space could require the solution of an integral equation for the determination of χ . (ii) Replace the Hydrogenic pre-factor $\Phi(\alpha, s)$ by the analytical HF wave function. (iii) Combine the expansion of the variational space with the improvement of the pre-factor. (iv) Employ a different analytical form for the wave function functional.

As noted above, it is also possible to search over all functions $\chi(q, s)$ such that the functional $\psi[\chi]$ leads to a physical property of interest. For example, let us consider the expectation of $W = r_1 + r_2$: $\langle r_1 + r_2 \rangle = \langle s \rangle = s_0 + \Delta s$. Here s_0 is the expectation from the pre-factor $\Phi(s)$. Assuming Δs known from experiment or some accurate calculation, and if a wave function functional $\psi[\chi]$ of the form in the above calculation is employed, then two distinct χ 's such that $\langle \psi[\chi] | s | \psi[\chi] \rangle = s_0 + \Delta s$ can be obtained by solution of the quadratic equation $a(q, s)\chi(q, s)^2 + 2b(q, s)\chi(q, s) + [c(q, s) - A] = 0$, where the constant $A = 2\Delta s/\alpha^4$, and where the coefficients a, b and c are the same as in Eq.(8-10). With the functionals $\psi[\chi_1]$ and $\psi[\chi_2]$ thus determined, the energy could then be obtained by minimization of the functional $I[\psi[\chi]]$ with respect to the parameters α and q . In this manner, the two wave function functionals would reproduce both the size of the atom exactly *and* the energy accurately.

For completeness we note that the concept of constrained search in the present work differs from that[7] within density functional theory (DFT). The idea underlying DFT is based on the first Hohenberg-Kohn theorem[8] according to which the wave function Ψ is a functional of the *ground* state density $\rho(\mathbf{r})$: $\Psi = \Psi[\rho]$. Thus the energy is a unique functional of $\rho(\mathbf{r})$: $E = E[\rho]$. The *in principle* constrained search within DFT is as follows. One first searches for the infimum of the $\langle \hat{H} \rangle$

TABLE I: Rigorous upper bounds to the ground state energy of the Helium atom in (a.u) for different wave functions (WF). The satisfaction of the virial theorem, and the expectation values of $W = r_1 + r_2$ are also quoted.

WF	Parameters	Energy	$-V/T$	$\langle W \rangle$
Φ	$\alpha = 1.6875$	-2.84766	2.0000	1.7778
$\psi[\chi_3]$	$\alpha = 1.6875 \ q = 0.581$	-2.89004	1.9678	1.7778
$\psi[\chi_4]$	$\alpha = 1.6875 \ q = 0.180$	-2.89061	1.9707	1.7778
$\psi[\chi_1]$	$\alpha = 1.6614 \ q = 0.5333$	-2.89072	1.9973	1.8057
$\psi[\chi_2]$	$\alpha = 1.6629 \ q = 0.17049$	-2.89122	1.9984	1.8041
<i>HF</i>	See Ref. 3	-2.86168	2.0000	1.8545
<i>CK</i>	See Ref. 5	-2.89007	1.9890	1.7848
<i>Pekeris</i>	See Ref. 4	-2.90372	2.0000	1.8589

over all antisymmetric, normalized, N -particle functions Ψ whose density is $\rho(\mathbf{r})$. One then searches over all such $\rho(\mathbf{r})$ to obtain the infimum of that expectation. The consecutive infima can be shown to be a minimum and equal to the ground state energy. In our work, the statement that the wave function is a functional of the functions χ is more general. The functions χ are not restricted to just being the density or functions of the density. And the constrained search over all functions χ is such that the wave function leads to an *arbitrary* property of interest. The energy is subsequently obtained by variational minimization.

In addition to its use in Schrödinger theory, the wave function functional $\psi[\chi]$ may also be employed within Quantal density functional theory (Q-DFT)[9]. In Q-DFT, as in traditional Kohn-Sham (KS) DFT[10], the system of electrons is transformed into one of *noninteracting* fermions such that the same ground state density, energy and ionization potential are obtained. However, in contrast to KS-DFT which is in terms of energy functionals and functional derivatives, the Q-DFT framework involves fields whose quantal sources are expectations of Hermitian operators taken with respect to the wave function. Thus, an approximate wave function functional of the form of Eq.(2) can be employed in this theory with the Slater determinantal pre-factor determined *self-consistently*. The wave function functional $\psi[\chi = f(\rho)]$ could also be used. Within Q-DFT, the corresponding energy is a rigorous upper bound.

We also mention the work of Colle and Salvetti(CS)[11] who suggested a wave function functional of the density. The CS wave function is similar to Eq.(2) except that the pre-factor is the HF wave function and the correlation term is $f(\mathbf{r}_1, \mathbf{r}_2) = e^{-\beta^2 r^2} [1 - \chi(\mathbf{R})(1 + r/2)]$, with $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, $\mathbf{R} = \mathbf{r}_1 + \mathbf{r}_2$, $\beta = q[\rho^{HF}(\mathbf{R})]^{1/3}$. CS further assumed that the corresponding Dirac density matrix was that due to the pre-factor. The

function $\chi(\mathbf{R})$ was to be determined by requiring that the correction to the HF Dirac density matrix due to the correlation factor $f(\mathbf{r}_1, \mathbf{r}_2)$ vanishes. Since the HF Dirac density matrix cannot be the same as the exact interacting system density matrix, this is not an exact constraint. Moreover, they did not satisfy this condition, and instead approximated the function $\chi(\mathbf{R})$ by $\chi(\mathbf{R}) = \sqrt{\pi}\beta/(1 + \sqrt{\pi}\beta)$. A consequence of this was that the resulting wave function was not normalized[5, 12, 13]. There is also no discussion in this work of the general concept of wave function functionals or the idea of constrained search to obtain them.

In conclusion, we have proposed the idea of expanding the space of variations beyond that of standard variational calculations by considering the wave function to be a functional instead of a function, a functional of the functions χ . A *constrained search* is performed over the functions χ such that the wave function satisfies a constraint or leads to a physical observable. A rigorous upper bound to the energy is then obtained by variational minimization with respect to any parameters in the wave function functional. The construction of such a constrained-search-variational wave function functional for the ground state of the Helium atom where the search is over the *entire* requisite subspace is explicitly demonstrated.

This work was supported in part by the Research Foundation of CUNY. L. M. was supported in part by NSF through CREST, and by a ‘‘Research Centers in Minority Institutions’’ award, RR-03037, from the National Center for Research Resources, National Institutes of Health.

-
- [1] E. A. Hylleraas, Z. Physik, **48**, 469 (1928).
 - [2] X.-Y. Pan and V. Sahni, J. Chem. Phys. **119**, 7083(2003).
 - [3] C. F. Fischer, *The Hartree-Fock Method For Atoms*, John Wiley and Sons, 1977.
 - [4] C. L. Pekeris, Phys. Rev. **115**, 1216 (1959).
 - [5] S. Caratzoulas and P. J. Knowles, Mol. Phys. **98**, 1811 (2000).
 - [6] J. Goodisman and W. Klemperer, J. Chem. Phys. **38**, 721(1963).
 - [7] M. Levy, *Proc. Natl. Acad. Sci. USA*, **76**,6062 (1979); E. H. Lieb, Int. J. Quantum Chem. **24**, 243 (1983).
 - [8] P. Hohenberg and W. Kohn, Phys. Rev. **136B**, 864 (1964).
 - [9] V. Sahni, *Quantal Density Functional Theory*, Springer Verlag (2004); V. Sahni and X.-Y. Pan, Phys. Rev. Lett. **90**, 123001(2003).
 - [10] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
 - [11] R. Colle and O. Salvetti, Theor. Chim. Acta **37**, 329 (1975).
 - [12] R. Singh et al, Phys. Rev. A **60**, 4135 (1999).
 - [13] J. Tao et al, Phys. Rev. A **63**, 032513(2001).

